

# Using adult mosquitoes to transfer insecticides to *Aedes aegypti* larval habitats

Gregor J. Devine<sup>a,1</sup>, Elvira Zamora Perea<sup>b</sup>, Gerry F. Killeen<sup>c,d</sup>, Jeffrey D. Stancil<sup>e,2</sup>, Suzanne J. Clark<sup>a</sup>, and Amy C. Morrison<sup>f</sup>

<sup>a</sup>Department of Plant and Invertebrate Ecology, Rothamsted Research, Harpenden AL5 2JQ, United Kingdom; <sup>b</sup>Laboratorio de Salud Publica, Iquitos, Peru; <sup>c</sup>Ifakara Health Institute, Dar es Salaam, United Republic of Tanzania; <sup>d</sup>Vector Group, Liverpool School of Tropical Medicine, Liverpool L3 5QA, United Kingdom; <sup>e</sup>Naval Medical Research Center Detachment, United States Navy, Lima, Peru; and <sup>f</sup>Department of Entomology, University of California, Davis, CA 95616

Edited by Barry J. Beaty, Colorado State University, Fort Collins, CO, and approved April 14, 2009 (received for review February 7, 2009)

Vector control is a key means of combating mosquito-borne diseases and the only tool available for tackling the transmission of dengue, a disease for which no vaccine, prophylaxis, or therapeutic currently exists. The most effective mosquito control methods include a variety of insecticidal tools that target adults or juveniles. Their successful implementation depends on impacting the largest proportion of the vector population possible. We demonstrate a control strategy that dramatically improves the efficiency with which high coverage of aquatic mosquito habitats can be achieved. The method exploits adult mosquitoes as vehicles of insecticide transfer by harnessing their fundamental behaviors to disseminate a juvenile hormone analogue (JHA) between resting and oviposition sites. A series of field trials undertaken in an Amazon city (Iquitos, Peru) showed that the placement of JHA dissemination stations in just 3–5% of the available resting area resulted in almost complete coverage of sentinel aquatic habitats. More than control mortality occurred in 95–100% of the larval cohorts of *Aedes aegypti* developing at those sites. Overall reductions in adult emergence of 42–98% were achieved during the trials. A deterministic simulation model predicts amplifications in coverage consistent with our observations and highlights the importance of the residual activity of the insecticide for this technique.

dengue | innovation | vector control | auto-dissemination | juvenile hormone analogue

Globally, 50 million dengue infections annually result in 500,000 cases of dengue hemorrhagic fever and 22,000 deaths (ref. 1 and [www.who.int/mediacentre/factsheets/fs117/en/print.html](http://www.who.int/mediacentre/factsheets/fs117/en/print.html)). *Aedes aegypti* (Linnaeus) transmits the bulk of dengue infections (2), and vector control is the only means of combating this disease for which no vaccine, prophylaxis, or therapeutic currently exists. The most effective means of controlling mosquito vectors of disease are insecticidal and include the use of adulticides as space sprays or indoor residual applications, insecticide-treated materials (ITMs) such as curtains and bed nets, and the application of larvicides to aquatic habitats (refs. 3–5 and [http://whqlibdoc.who.int/trs/WHO\\_TRS\\_857.pdf](http://whqlibdoc.who.int/trs/WHO_TRS_857.pdf)). These tools may be augmented by source reduction campaigns targeted at mosquito breeding sites (6, 7). The primary challenge for the effective implementation of any of these measures is in realizing sufficient coverage of the insect population given local constraints on financial and human resources (3, 7–9).

The application of adulticides and the use of treated bed nets can have a powerful impact on the abundance of mosquito vectors (10, 11) and disease transmission (12, 13) because the host-seeking and resting behaviors of the vector ensure a number of potentially lethal interactions with insecticide-treated surfaces during those parts of the lifecycle when pathogens are acquired, incubated, and transmitted. Mosquito density, longevity, and feeding success, which are some of the key determinants of vectorial capacity and disease transmission (14, 15), are all

affected. The efficacy of these tools, however, against many disease vectors, is often constrained by the difficulty in achieving sufficiently high coverage of resting surfaces, sleeping spaces, or adult vectors (7–9, 16). Aquatic habitat management can also contribute to decreasing transmission of mosquito-borne diseases (17, 18) but is often considered inferior to adulticiding and ITMs because it does not impact directly on the most important determinants of vectorial capacity. To exert a significant effect on transmission, aquatic habitat management methods depend on simply maximizing their impacts on adult mosquito density. At large or spatially complex scales this is challenging, because of uncertainty over the relative productivity of specific habitats and the consequent need to seek out, identify, and treat all potential sites (6, 7).

The strategy that we describe here exploits the innate behaviors of adult mosquitoes to effectively target a persistent juvenile hormone analogue (JHA) at their aquatic habitats. Adult females, exposed to JHA deposits at their resting sites, contaminate aquatic habitats and the larvae developing therein when they oviposit. The tiny doses of JHA that they transfer then interfere with the metamorphosis of those juvenile stages. We demonstrate, in theory and practice, that high coverage of aquatic habitats with a JHA is possible through the treatment of only a small proportion of the adult resting area. This has a marked impact on the emergence of adults from contaminated sites. The impetus for our field demonstrations was given by some highly artificial, laboratory-based explorations of the insecticide-transfer principle (19–21) and by a further characterization of the technique's potential using large cages and free-flying mosquitoes (*SI Text* and *Figs. S1 and S2*).

## Results

In 3 separate trials, undertaken in each of 2 sites in a public cemetery in the Amazon (Iquitos, Peru), we examined the impact of deploying 10 JHA “dissemination stations” on the productivity of 40 uncontaminated sentinel oviposition sites (Fig. 1). Each of these sentinel habitats contained a cohort of 25 uncontaminated third-instar *A. aegypti* larvae. When no JHA was deployed, the juvenile stages developing in the sentinel sites exhibited average mortalities of 8% (site A) and 7% (site B). During the postdeployment phase, mortality increased to 84% at site A (all dates combined;  $F = 78.9$ ,  $P < 0.001$ ) and 49% at site

Author contributions: G.J.D., G.F.K., J.D.S., and A.C.M. designed research; G.J.D. and E.Z.P. performed research; G.F.K. contributed new reagents/analytic tools; S.J.C. analyzed data; and G.J.D. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.


Freely available online through the PNAS open access option.

<sup>1</sup>To whom correspondence should be addressed. E-mail: [greg.devine@bbsrc.ac.uk](mailto:greg.devine@bbsrc.ac.uk).

<sup>2</sup>Present address: Joseph H. Pendleton Marine Corps Base Camp, San Diego, CA 92055.


This article contains supporting information online at [www.pnas.org/cgi/content/full/0901369106/DCSupplemental](http://www.pnas.org/cgi/content/full/0901369106/DCSupplemental).

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>APR 2009</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2009 to 00-00-2009</b>	
4. TITLE AND SUBTITLE <b>Using adult mosquitoes to transfer insecticides to Aedes aegypti larval habitats</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>University of California, Department of Entomology, Davis, CA, 95616</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>5</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			



**Fig. 1.** Schematic of experimental design in the Iquitos public cemetery. (A) Schematic of avenues and tombs in cemetery (not to scale). (B) Detail of tomb wall. (C) JHA dissemination station in a tomb. (D) Positioning of dissemination stations and sentinel sites (not to scale). Gray circles indicate sentinel sites with larval cohorts ( $n = 40$ ). Black circles indicate dissemination stations treated with JHA ( $n = 10$ ).

B (all dates combined;  $F = 55.7$ ,  $P < 0.002$ ). The maximum mortality seen in individual trials was 98% and 59% at sites A and B, respectively (Fig. 2). The effects of the JHA were most



**Fig. 2.** Effect of the presence of JHA-treated dissemination stations on the mortality of juvenile cohorts developing in sentinel sites (mean  $\pm$  95% confidence limits). Schematic shows how adults transfer JHA to the sentinel sites.


apparent on the nonfeeding pupal stage. Pupae accounted for 91.5% ( $n = 3552$ ) of posttreatment mortality but only 32.5% ( $n = 156$ ) of pretreatment mortality ( $F = 264$ ,  $P < 0.0001$ ).

The dissemination of the insecticide was extremely efficient. By placing a JHA-treated station in just 3% and 5% of the tombs available at sites A and B, respectively, we exerted a lethal effect on almost every sentinel site. After JHA deployment, only 3 sentinel sites (1 at site A and 2 at site B) exhibited mortality rates equal to or lower than those noted during the predeployment period (Fig. 3). This result suggests that the vast majority of sentinel sites ( $\geq 95\%$ ) in any trial were visited by contaminated mosquitoes. Distances between dissemination stations and sentinel sites were small and, at these scales, the mortality observed in individual aquatic habitats was not related to their distance from the 10 dissemination stations (Fig. 3).

A simple deterministic simulation model was used to demonstrate how the persistence of the JHA and/or multiple contaminations by disseminating adults can amplify the effective coverage of aquatic habitats. Further details of the model assumptions and explanations of its parameters are provided in *SI Text* and *Table S1*. The model proposes that the relationship between the coverage of adult resting sites ( $C_r$ ) and the larval habitats that the JHA is disseminated to ( $C_h$ ) can be crudely described as a simple exponential function of the duration for which habitats remain unproductive after contamination ( $U$ ), the number of ovipositions by the vector population ( $O$ ) relative to the number of habitats ( $H$ ), and the mean number of contaminated ovipositions required to render a single habitat unproductive ( $\Omega$ ):

$$C_h = 1 - \exp(-C_r U [O/H\Omega]).$$

Fig. 4A illustrates that, by using 1/20th of the available resting sites ( $C_r = 0.05$ ) to disseminate the insecticide, more than half of the larval habitats ( $C_h > 0.5$ ) can be affected (an amplification in coverage by a factor of  $>10$ ) given the following criteria: (i) aquatic habitats are rendered unproductive for at least 1 week,  $U \geq 7$  days; (ii) mosquito abundance or habitat availability is such that aquatic habitats are oviposited in more than once per 24 h,  $O/H \geq 2$ ; and (iii) only 1 contamination event is necessary to render a habitat unproductive,  $\Omega = 1$ . Increasing the persistence of the insecticide ( $U \geq 14$ ) and the number of oviposition events in each habitat ( $O/H \geq 5$ ) leads to almost complete habitat coverage ( $C_h > 0.95$ ; Fig. 4A). Fig. 4B uses the same model to illustrate how the persistence of the insecticide ( $U$ ) is the key to



**Fig. 3.** Postdeployment mortality in individual sentinel sites during the field trials. Points marked with \* denote the only 3 sentinel sites that showed  $\leq$  control mortality (i.e.,  $\leq$  the higher 95% CI displayed in Fig. 2). Simple regressions of sentinel site mortality against average distances from JHA dissemination stations are shown for trials 1 (black circles and line), 2 (gray circles and line), and 3 (white circles and dashed lines).

amplifying coverage between resting exposure ( $C_r$ ) and aquatic habitats ( $C_h$ ). These outcomes approximate the results of our field demonstrations in which the placement of dissemination traps treated with a persistent JHA in  $<5\%$  of available tombs ( $C_r < 0.05$ ) for 12-day periods resulted in the almost complete coverage of sentinel sites ( $C_h \geq 0.95$ ) with the insecticide.

## Discussion


The strength and impact of the strategy that we have described derives from the amplification in JHA coverage that results from the repeated contamination of adult mosquitoes at their resting

sites and the persistence of the insecticide. This efficient dissemination process is facilitated by the cyclical nature of mosquito feeding, resting, and oviposition behaviors.

Our trials showed that the wild mosquito population moved the JHA around the cemetery very effectively. Almost all of the sentinel sites were affected, despite the fact that the various avenues and niches of the cemetery presented myriad resting and oviposition opportunities. Moreover, there was no loss of impact on the sentinel site cohorts with increasing distance from the JHA dissemination stations. This suggests that the JHA was being disseminated beyond the boundaries of our study site and that we might reduce resting site coverage further while maintaining similar or greater impacts on larval habitats, at larger scales.

The JHA that we used (pyriproxyfen) does not interfere with the fundamental behaviors that we are exploiting because it is neither lethal nor repellent to adults (21). It is the act of oviposition that contaminates the aquatic habitat, so the technique explicitly and precisely targets the mosquitoes' preferred larval development sites. This may help overcome one of the most important constraints on the successful application of larvicidal or pupacidal interventions, the inefficient waste of expensive insecticides and human resources on treating inappropriate or cryptic oviposition sites.

Previous studies have shown that *A. aegypti* oviposit in a number of different habitats over the course of their gonotrophic cycle (22, 23) thus permitting a number of transfer events between resting and oviposition sites. In urban Iquitos, water volumes of 3–15 L account for the majority of *Aedes*-positive containers and the greatest pupal abundance (24). Positive containers tend to be those that are unmanaged (i.e., passively collected water is left standing for some days), which may facilitate the accumulation of lethal JHA concentrations through successive contamination events. Once lethal doses are achieved, pyriproxyfen can render domestic water storage containers unproductive for months rather than weeks (21). Adult *A. aegypti* mosquitoes are well suited for exploitation using this transfer technique because their resting sites have been well-described [e.g., dark spaces in houses (25, 26)] and appropriate dissemination traps are therefore simple to design and distribute. The concentration of pyriproxyfen that prevents adult emergence from local populations of third-instar *A. aegypti* larvae ( $LC_{50}$ ) is 0.012 parts per billion (ppb) (21), which is equivalent to the transfer of  $\approx 0.4 \mu\text{g}$  of JHA dust to a 200-mL aquatic sentinel site; just 1/1,000th of the dry weight of an *A. aegypti* adult (27). Scanning electron microscopy of contaminated mosquitoes helps visualize the ease with which such tiny doses can be picked up on the tarsi of resting mosquitoes (Fig. 5). Adult *Aedes*



**Fig. 4.** Deterministic simulation model outcomes. (A) Resting site coverage ( $C_r$ ) can be amplified by insecticide persistence ( $U$ ) and the number of ovipositions per habitat ( $O/H$ ) to achieve high habitat coverage ( $C_h$ ). (B) Under stable conditions of contamination ( $O/H = 1$ ) the persistence of the insecticide ( $U$ ) is the key to achieving high habitat coverage.





equivalent of 5 g of pyriproxyfen/m<sup>2</sup> (Sumilarv 0.5G; Sumitomo Chemical Corporation; a 0.5% granular formulation) pulverized to the consistency of talcum powder. The water in these stations served to dampen the cloth lining and ensure that the pyriproxyfen remained stuck to the cloth and available to resting mosquitoes.

In addition to the 10 dissemination stations, we distributed 40 sentinel oviposition sites among the lower 3 rows of tombs in each wall. These consisted of 1-L disposable containers holding 200 mL of water and 25 uncontaminated, laboratory-reared, late third-instar *A. aegypti* larvae. All sentinel sites were between 1.05 and 37.5 m from each of the 10 dissemination stations deployed in those avenues (Fig. 1D). Because the contamination of any single sentinel site could result from the transfer of JHA from any or all of the 10 dissemination stations, correlations between sentinel site mortality and proximity to dissemination stations are presented in terms of an average cumulative distance (Fig. 3).

During the trials, each of the 40 sentinel sites and their attendant larval cohorts were monitored every day. Dead larvae and pupae were counted and discarded. Live larvae were left in the cemetery to develop further but live pupae were counted, removed by pipette, and placed in a disposable cup (a separate cup for each artificial oviposition site) containing uncontaminated water. These cups were covered with gauze lids and taken to the laboratory where they were maintained at  $27 \pm 3$  °C. This process ensured that there was no release of mosquitoes into the cemetery. This procedure continued until none of the original cohort remained in the sentinel sites. In all cases, this required 12 days or less, although observations of live pupae, removed from

the site, continued in the laboratory. These laboratory-maintained pupae were monitored daily until they emerged as adults or died. Thus, for each sentinel cohort, we derived cumulative totals of dead larvae, dead pupae, emerged adults, and overall mortality. Any discrepancy between the final totals and the 25 larvae originally placed in each pot (i.e., missing larvae) was added to the mortality total (cadavers and weak individuals often disappear as they are scavenged by older instar larvae). Natural populations of larvae, resulting from oviposition by wild adults, were periodically removed from the pots before they reached third instar, so that they could not be confused with the late-instar, laboratory-reared cohort that was being monitored.

For the field tests, all proportional data were transformed [ $\arcsin(\sqrt{p})$ ] for analysis by ANOVA and t test. Data are presented as back transformed means and 95% confidence limits.

**ACKNOWLEDGMENTS.** The Directors of the local Dirección de Salud, the Laboratorio de Salud Pública, and the Comité de Investigaciones in Peru gave written consent for the trials and associated protocols. Sra. Rosa Ochoa de Zamora, president of the Beneficencia Pública de Iquitos, allowed us access to the public cemetery. We thank Ernesto Curto, the director of the Laboratorio de Salud Pública, Iquitos for the use of the entomology facilities and Wagner Orellana Rios and Andreas Rojas Estrella for expert technical assistance. G.J.D. was partially supported by U.S. Navy Works Unit Number 62787A 870 U8517 and a Deployed War Fighter Program (2005–2007) grant (to G.J.D., J.D.S., and A.C.M.) administered by the U.S. Armed Forces Pest Management Board. G.F.K. was supported by Research Career Development Fellowship 076806 from the Wellcome Trust.

- WHO (2008) *Dengue and Dengue Haemorrhagic Fever* (WHO, Geneva), WHO Fact Sheet 117.
- Phillips M (2008) Dengue reborn: Widespread resurgence of a resilient vector. *Environ Health Perspect* 116:A382–A388.
- Anonymous (1997) The feasibility of eradicating *Aedes aegypti* in the Americas. *Rev Panam Salud Publica Pan Am J Public Health* 1:68–72.
- Ritchie S, et al. (2002) Dengue control in North Queensland, Australia: Case recognition and selective indoor residual spraying. *Dengue Bull* 26:7–13.
- WHO (1995) *Vector Control for Malaria and Other Mosquito Borne Diseases* (WHO, Geneva), WHO Technical Report Series.
- Fillinger U, Sonye G, Killeen GF, Knols BGJ, Becker N (2004) The practical importance of permanent and semipermanent habitats for controlling aquatic stages of *Anopheles gambiae* sensu lato mosquitoes: Operational observations from a rural town in western Kenya. *Trop Med Int Health* 9:1274–1289.
- Morrison AC, Zielinski-Gutierrez E, Scott TW, Rosenberg R (2008) Defining challenges and proposing solutions for control of the virus vector *Aedes aegypti*. *PLoS Med* 5:362–366.
- Curtis CF, Mnzava AEP (2000) Comparison of house spraying and insecticide-treated nets for malaria control. *Bull WHO* 78:1374–1491.
- Killeen GF, et al. (2007) Preventing childhood malaria in Africa by protecting adults from mosquitoes with insecticide-treated nets. *PLoS Med* 4:1246–1258.
- Kroeger A, et al. (2006) Effective control of dengue vectors with curtains and water container covers treated with insecticide in Mexico and Venezuela: Cluster randomized trials. *Br Med J* 332:1247–1252.
- Lindblade KA, et al. (2006) Impact of sustained use of insecticide-treated bednets on malaria vector species distribution and culicine mosquitoes. *J Med Entomol* 43:428–432.
- Lengeler C (2004) Insecticide-treated bed nets and curtains for preventing malaria. *Cochrane Database Syst Rev* 2:CD000363.
- Lenhart A, et al. (2008) Insecticide-treated bednets to control dengue vectors: Preliminary evidence from a controlled trial in Haiti. *Trop Med Int Health* 13:56–67.
- Dye C (1984) Models for the population dynamics of the yellow-fever mosquito, *Aedes aegypti*. *J Anim Ecol* 53:247–268.
- Macdonald G (1957) *The Epidemiology and Control of Malaria* (Oxford Univ Press, London).
- Reiter P (2007) Oviposition, dispersal, and survival in *Aedes aegypti*: Implications for the efficacy of control strategies. *Vector Borne Zoonotic Dis* 7:261–273.
- Killeen GF, Fillinger U, Kiche I, Gouagna LC, Knols BGJ (2002) Eradication of *Anopheles gambiae* from Brazil: Lessons for malaria control in Africa? *Lancet Infect Dis* 2:618–627.
- Sanchez L, et al. (2006) *Aedes aegypti* larval indices and risk for dengue epidemics. *Emerging Infect Dis* 12:800–806.
- Chism BD, Apperson CS (2003) Horizontal transfer of the insect growth regulator pyriproxyfen to larval microcosms by gravid *Aedes albopictus* and *Ochlerotatus triseriatus* mosquitoes in the laboratory. *Med Vet Entomol* 17:211–220.
- Itoh T, et al. (1994) Utilization of blood-fed females of *Aedes aegypti* as a vehicle for the transfer of the insect growth regulator pyriproxyfen to larval habitats. *J Am Mosq Cont Assoc* 10:344–347.
- Sihuinchu M, et al. (2005) Potential use of pyriproxyfen for control of *Aedes aegypti* (Diptera: Culicidae) in Iquitos, Peru. *J Med Entomol* 42:620–630.
- Apostol BL, Black WC, Reiter P, Miller BR (1994) Use of randomly amplified polymorphic DNA amplified by polymerase chain-reaction markers to estimate the number of *Aedes aegypti* families at oviposition sites in San Juan, Puerto Rico. *Am J Trop Med Hyg* 51:89–97.
- Reiter P, Amador M, Anderson R, Clark G (1995) Short report: Dispersal of *Aedes aegypti* in an urban area after blood feeding as demonstrated by rubidium-marked eggs. *Am J Trop Med Hyg* 52:177–179.
- Morrison AC, et al. (2004) Temporal and geographic patterns of *Aedes aegypti* (Diptera: Culicidae) production in Iquitos, Peru. *J Med Entomol* 41:1123–1142.
- Edman J, Kittayapong P, Linthicum K, Scott T (1997) Attractant resting boxes for rapid collection and surveillance of *Aedes aegypti* (L) inside houses. *J Am Mosq Control Assoc* 13:24–27.
- Perich MJ, Davila G, Turner A, Garcia A, Nelson M (2000) Behavior of resting *Aedes aegypti* (Culicidae: Diptera) and its relation to ultra-low volume adulticide efficacy in Panama City, Panama. *J Med Entomol* 37:541–546.
- Nasci RS (1990) Relationship of wing length to adult dry-weight in several mosquito species (Diptera, Culicidae). *J Med Entomol* 27:716–719.
- Agudelo-Silva F, Spielman A (1984) Paradoxical effects of simulated larviciding on production of adult mosquitoes. *Am J Trop Med Hyg* 33:1267–1269.
- Wilson ML, Agudelo-Silva F, Spielman A (1990) Increased abundance size and longevity of food-deprived mosquito populations exposed to a fungal larvicide. *Am J Trop Med Hyg* 43:551–556.
- Ivanaga K, Kanda T (1988) The effects of a juvenile-hormone active oxime ether compound on the metamorphosis and reproduction of an anopheline vector, *Anopheles balabacensis* (Diptera, Culicidae). *Appl Entomol Zool* 23:186–193.
- WHO (2004) *Pyriproxyfen in Drinking Water. Background Document for Development of WHO Guidelines for Drinking-Water Quality* (WHO, Geneva).
- Lazcano JAB, et al. (2006) Ecological factors linked to the presence of *Aedes aegypti* larvae in highly infested areas of Playa, a municipality belonging to Ciudad de La Habana, Cuba. *Rev Panam Salud Publica Pan Am J Public Health* 19:379–384.
- Harrington LC, et al. (2005) Dispersal of the dengue vector *Aedes aegypti* within and between rural communities. *Am J Trop Med Hyg* 72:209–220.
- Gillies M, de Meillon B (1968) *The Anophelini of Africa south of the Sahara (Ethiopian zoogeographical region)* (South African Institute of Medical Research, Johannesburg, South Africa), 2nd Ed.
- Charlwood JD, Vij R, Billingsley PF (2000) Dry season refugia of malaria-transmitting mosquitoes in a dry savannah zone of east Africa. *Am J Trop Med Hyg* 62:726–732.
- Minakawa N, Sonye G, Yan GY (2005) Relationships between occurrence of *Anopheles gambiae* s.l. (Diptera: Culicidae) and size and stability of larval habitats. *J Med Entomol* 42:295–300.
- Killeen GF, et al. (2000) The potential impact of integrated malaria transmission control on entomologic inoculation rate in highly endemic areas. *Am J Trop Med Hyg* 62:545–551.
- Invest JF, Lucas JR (2008) Pyriproxyfen as a mosquito larvicide. *Sixth International Conference on Urban Pests*, eds Robinson WH, Bajomi D (OOK-Press Kft, Veszprém, Hungary), pp 239–245.
- Nauen R (2007) Insecticide resistance in disease vectors of public health importance. *Pest Manag Sci* 63:628–633.
- Morrison AC, et al. (2006) *Aedes aegypti* (Diptera: Culicidae) production from nonresidential sites in the Amazonian city of Iquitos, Peru. *Ann Trop Med Parasitol* 100:573–586.